

Chiral copper complex and production processes thereof and using the same

Field of the invention

The present invention relates to a chiral copper complex, the
5 production process thereof using chiral salicylidene ligands and asymmetric
synthesis of cyclopropane-carboxylic acid compound using the complex as a
catalyst.

Background of the Invention

10 As a process for producing an optically active cyclopropanecarboxylic
acid ester derivative, there have been known a process of reacting an olefin
with a diazoacetic acid ester in the presence of a certain
salicylideneaminoalcohol copper complex catalyst (JP-A 59-225194).


15 Summary of the Invention

According to the present invention, an industrially suitable chiral
copper complex comprising an optically active salicylidenaminoalcohol
compound can be readily produced. Said complex is more stable to an oxide
that can be contained in a cyclopropanation reaction system, reducing
20 adverse effects of such an oxide and said salicylidenaminoalcohol compound
can be recovered after said reaction in an improved yield.

The present invention provides:

1. an optically active salicylidenaminoalcohol compound of formula
(1):

25



wherein R₁ represents

an alkyl group which may be substituted with a group selected from
an alkoxy group, an aralkyloxy group, an aryloxy group and cycloalkoxy
30 group,

an aralkyl, aryl or cycloalkyl group all of which may be substituted

with a group selected from an alkyl group, an alkoxy group, an aralkyloxy group, an aryloxy group, and a cycloalkoxy group,

R_2 represents

an alkyl group, a cycloalkyl group, or

5 an aralkyl or phenyl group which may be substituted with a group selected from an alkyl group, an alkoxy group, an aralkyloxy group, an aryloxy group and a cycloalkoxy group,

when X_1 represents a nitro, X_2 is a hydrogen atom,

when X_1 represents a chlorine atom, X_2 is a chlorine atom, and


10 when X_1 is a hydrogen atom, X_2 is a fluorine atom, and

the carbon atom denoted by " * " is an asymmetric carbon atom having either an S or R configuration,


2. a process for producing an optically active salicylideneaminoalcohol compound as defined above,

15 which comprises reacting

an optically active amino alcohol of formula (2):




wherein R_1 , R_2 and " * " have the same meanings as defined above, with
20 a 2-hydroxybenzaldehyde derivative of formula (3):



wherein when X_1 and X_2 are the same as defined above,

3. a chiral copper complex prepared from a mono-valent or di-valent
25 copper compound and an optically active salicylideneaminoalcohol compound (1) as defined above,

4. a method for producing a chiral copper complex of formula (1):



wherein R_1 and R_2 are the same or different and independently represent an alkyl group, an aralkyl group, a phenyl group, a 2-methoxyphenyl group,
 5 a 2-tert-butoxy-5-tert-butylphenyl group, or a 2-octyloxy-5-tert-butylphenyl group;

when X_1 represents a nitro group, X_2 is a hydrogen atom,


when X_1 represents a chlorine atom, X_2 is a chlorine atom, and

when X_1 represents a hydrogen atom, X_2 is a fluorine atom

10 the carbon atom denoted by " * " is an asymmetric carbon atom having either an S or R configuration,


which comprises contacting a di-valent copper compound, in an inert organic solvent, with a chiral salicylideneaminoalcohol compound of formula (1):

15




wherein R_1 , R_2 , X_1 , X_2 and " * " respectively have the same meaning as defined above, and

5. a method for producing an optically active
 20 cyclopropane-carboxylic acid ester of formula (4):



wherein R_3 , R_4 , R_5 , R_6 and R_7 are as defined below,

which comprises reacting a prochiral olefin of formula (5):



wherein R_3 , R_4 , R_5 and R_6 are as defined below, with a diazoacetic acid ester of formula (6):



wherein R_7 is as defined below, in the presence of a chiral copper complex as defined in item 3 or 4,

wherein R_3 , R_4 , R_5 and R_6 independently represent

a hydrogen atom,

a halogen atom,

a (C1-C8)alkyl group which may be substituted with a halogen atom

or a lower alkoxy group,

a (C4-C8)cycloalkyl group,

an aryl group which may be substituted with a halogen atom or a lower alkoxy group,

an alkoxy group,

R_3 and R_4 , or R_5 and R_6 may be bonded at their terminals to form an alkylene group having 2-4 carbon atoms, and

one of R_3 , R_4 , R_5 and R_6 groups represents an alkenyl group which may be substituted with a halogen atom, an alkoxy group or an alkoxy carbonyl group, of which alkoxy may be substituted with a halogen atom or atoms,

provided that when R_3 and R_5 are the same, R_4 and R_6 are not the same, and

R_7 represents an alkyl group having 1 to 8 carbon atoms,

a cycloalkyl group which may be optionally substituted with a lower alkyl group,

a benzyl group which may be optionally substituted with a lower alkyl group, a lower alkoxy group, a phenoxy group or a halogen atom,

a phenyl group which may be optionally substituted with a lower alkyl group, a lower alkoxy group or a phenoxy group,

Detailed Description

First, a description will be made to the optically active salicylideneaminoalcohol compound of formula (1) as defined above.

5 Examples of the alkyl group which may be substituted with a group selected from an alkoxy group, an aralkyloxy group, an aryloxy group, and a cycloalkoxy group, represented by R_1 , include

 a (C1-C8)alkyl group (e.g., a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, n-pentyl, n-hexyl, n-pentyl, n-octyl, n-nonyl, or
10 n-decyl group) which may be substituted with a group selected from a (C1-C4)alkoxy group (e.g., a methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, i-butoxy, sec-butoxy, or t-butoxy group),

 a (C7-C11)aralkyloxy group (e.g., a benzyloxy, or naphthylmethyloxy group),

15 a (C6-C11)aryloxy group (e.g., a phenoxy, or naphthoxy group) and

 a (C4-C6)cycloalkoxy group (e.g., cyclobutyloxy, cyclopentyloxy, cyclohexyloxy group).

 Examples of the aralkyl group, the aryl group, and the cycloalkyl group, all of which may be substituted with a group selected from an alkyl group, an alkoxy group, an aralkyloxy group, an aryloxy group, and a
20 cycloalkoxy group include

 a (C7-C11)aralkyl group (e.g., a benzyl, or naphthylmethyl group),

 a (C6-C10)aryl group (e.g., a phenyl, or naphthyl group),

25 a (C4-C6)cycloalkyl group (e.g., a cyclobutyl, cyclopentyl, or cyclohexyl group), all of which may be substituted with a group as specified above.

 The alkyl group represented by R_2 has the same meanings as defined above for the (C1-C8)alkyl group.

 The cycloalkyl group, and the aralkyl or phenyl group which may be substituted with a group selected from an alkyl group, an alkoxy group, an
30 aralkyloxy group, an aryloxy group and a cycloalkoxy group represented by R_2 have the same meanings as defined above for R_1 .

 Among the optically active salicylideneaminoalcohol compound of formula (1), R_1 and R_2 are preferably an alkyl group (e.g. lower alkyl groups having 1 to 4 carbon atoms such as methyl group, ethyl group, n-propyl group,
35 isopropyl group, n-butyl group, tert-butyl group or the like), an aralkyl group (e.g., a benzyl group), an aryl group (e.g., a phenyl group, a 2-methoxyphenyl

group, a 2-tert-butoxy-5-tert-butylphenyl group or a 2-octyloxy-5-tert-butylphenyl group).

Specific examples the optically active salicylideneaminoalcohol compound of formula (1) include optically active

- 5 N-(3-fluorosallylidene)-2-amino-1,1-di(2-butoxy-5-tert-butylphenyl)-1-propanol,
N-(3-fluorosallylidene)-2-amino-1,1-di(2-octyloxy-5-tert-butylphenyl)-1-propanol,
N-(3-fluorosallylidene)-2-amino-1,1-di(2-butoxy-5-tert-butylphenyl)-3-phenyl-1-propanol,
N-(3-fluorosallylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol,
10 N-(3-fluorosallylidene)-2-amino-1,1-diphenyl-1-propanol,
N-(3-fluorosallylidene)-2-amino-1,1-di(2-benzyloxy-5-methylphenyl)-3-(4-isopropoxyphenyl)-1-propanol,
N-(3-fluorosallylidene)-2-amino-1,1-diphenyl-3-phenyl-1-propanol,
N-(3-fluorosallylidene)-2-amino-1,1-di(2-methoxyphenyl)-3-methyl-1-butanol,
15 N-(3,5-dichlorosallylidene)-2-amino-1,1-diphenyl-1-propanol,
N-(3,5-dichlorosallylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol
N-(3,5-dichlorosallylidene)-2-amino-1,1-di(5-tert-butyl-2-tert-butoxyphenyl)-3-phenyl-1-propanol,
N-(3,5-dichlorosallylidene)-2-amino-1,1-di(5-tert-butyl-2-octyloxyphenyl)-1-propanol
20 N-(3,5-dichlorosallylidene)-2-amino-1,1-di(2-methoxyphenyl)-3-methyl-1-butanol,
N-(5-nitrosallylidene)-2-amino-1,1-diphenyl-1-propanol,
N-(5-nitrosallylidene)-2-amino-1,1-diphenyl-3-phenyl-1-propanol,
25 N-(5-nitrosallylidene)-2-amino-1,1-di(2-butoxy-5-tert-butylphenyl)-3-phenyl-1-propanol,
N-(5-nitrosallylidene)-2-amino-1,1-di(2-benzyloxy-5-methylphenyl)-3-(4-isopropoxyphenyl)-1-propanol,
N-(5-nitrosallylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol,
30 N-(5-nitrosallylidene)-2-amino-1,1-di(2-tert-butyl-4-methylphenyl)-3-phenyl-1-propanol,
N-(5-nitrosallylidene)-2-amino-1,1-di(4-tert-butylphenyl)-1-propanol,
N-(5-nitrosallylidene)-2-amino-1,1-di(2-methoxyphenyl)-3-methyl-1-propanol,
N-(5-nitrosallylidene)-2-amino-1,1-di(5-tert-butyl-2-octyloxyphenyl)-1-propanol,
35 N-(5-nitrosallylidene)-2-amino-1,1-di(5-tert-butyl-2-tert-butoxyphenyl)-3-

phenyl-1-propanol,

N-(5-nitrosalicylidene)-2-amino-1,1-di(5-tert-butyl-2-tert-butoxyphenyl)-3-methyl-1-butanol and the like. Said optically active compounds have either an S configuration or R configuration with respect to the carbon atom
5 denoted by " * " in the formula above.

Next, a description will be made to the process for producing the optically active salicylideneaminoalcohol compound of formula (1) comprising reacting an optically active amino alcohol of formula (2) with a salicylaldehyde derivative of formula (3) as defined above.

10 The optically active amino alcohol compound of formula (2) to be used in this process include those having R₁ and R₂ groups as specified above and specific examples thereof include optically active

2-amino-1,1-diphenyl-1-propanol,

2-amino-1,1-di(2-methoxyphenyl)-1-propanol,

15 2-amino-1,1-di(2-isopropoxyphenyl)-1-propanol,

2-amino-1,1-di(2-butoxy-5-t-butylphenyl)-1-propanol,

2-amino-1,1-diphenyl-3-phenyl-1-propanol,

2-amino-1,1-di(2-methoxyphenyl)-3-phenyl-1-propanol,

2-amino-1,1-di(2-isopropoxyphenyl)-3-phenyl-1-propanol,

20 2-amino-1,1-di(2-butoxy-5-t-butylphenyl)-3-phenyl-1-propanol,

2-amino-1,1-di(2-methoxyphenyl)-3-phenyl-1-butanol,

2-amino-1,1-di(2-butoxy-5-t-butylphenyl)-1-propanol,

2-amino-1,1-di(2-octyloxy-5-t-butylphenyl)-1-propanol,

2-amino-1,1-di(2-butoxy-5-t-butylphenyl)-3-phenyl-1-propanol,

25 2-amino-1,1-di(2-methoxyphenyl)-1-propanol,

2-amino-1,1-di(2-benzyloxy-5-methylphenyl)-3-(4-isopropoxyphenyl)-1-propanol,

2-amino-1,1-di(2-methoxyphenyl)-3-methyl-1-butanol,

2-amino-1,1-di(2-t-butyl-4-methylphenyl)-3-phenyl-1-propanol,

30 2-amino-1,1-di(4-t-butylphenyl)-1-propanol,

2-amino-1,1-di(2-methoxyphenyl)-3-methyl-1-propanol and the like.

The reaction of the optically active amino alcohol (2) with the salicylaldehyde derivative (3) is usually conducted at 20 to 150°C, preferably 50 to 120°C.

35 Said reaction is usually conducted by contacting the optically active amino alcohol with the salicylaldehyde derivative (3) in an organic solvent,

examples of which include

an aromatic hydrocarbon solvent such as toluene, xylene or the like,
a halogenated hydrocarbon solvent such as chlorobenzene,
dichloromethane, dichloroethane or the like,

5 an alcohol solvent such as methanol or the like,
an aliphatic hydrocarbon such as hexane, heptane, cyclohexane or the
like,

an ether such as diethylether, methyl-t-butylether or the like, an
ester such as methyl acetate, ethyl acetate or the like, and a mixture thereof.

10 An amount thereof to be used is not particularly limited.

An amount of the salicylaldehyde derivative to be used is usually 1 to 2
moles, preferably 1 to 1.5 moles per mol of the optically active amino alcohol
of formula (2).

The salicylideneaminoalcohol of formula (1) is usually contacted with a
15 mon-valent or di-valent copper compound to produce a chiral copper complex
in a solvent.

Examples of the mono-valent or di-valent copper compound include

a copper salt of an organic carboxylic acid having 2 to 15 carbon
atoms such as copper acetate, copper naphthenate, copper octanoate and the
20 like, and

a copper salt such as copper chloride, copper bromide, copper nitrate,
copper sulfate, copper methanesulfonate, copper trifluoromethanesulfonate,
copper cyanate, or copper carbonate, and copper oxide, and a mixture thereof.

25 Examples of the solvent include a halogenated hydrocarbon such as
dichloromethane, dichloroethane, chloroform, carbon tetrachloride or the like,
and an aromatic hydrocarbon such as toluene, xylene or the like, an ether
such as diethylether, methyl-t-butylether or the like, an ester such as methyl
acetate, ethyl acetate or the like, and a mixture thereof. An amount thereof
to be used is not particularly limited and is usually 2 to 500 parts by weight
30 per 1 part by weight of the copper compound. Prochiral olefins of formula (6)
to be used in the next cyclopropanation step may also be used as a solvent.

An amount of the optically active salicylideneaminoalcohol compound
of formula (1) to be used is usually 0.5 to 5 moles, preferably 0.8 to 2.5 moles
per mol of the copper compound. The reaction temperature is usually 10 to
35 120°C, preferably 20 to 100°C.

After completion of the reaction, the resulting reaction mixture

containing the desired chiral copper complex may be used as it is in a solution form in the subsequent cyclopropanation reaction, or it may be washed with an aqueous alkali solution (e.g, aqueous solution of sodium hydrogencarbonate, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate or the like) and the evaporated to give the
5 desired chiral copper complex.

Alternatively the desired copper complex can be precipitated as a green colored crystal by cooling the reaction solution at a temperature of ranging from -50 to +30°C, preferably -10 to 20°C.

10 The copper complex may be also precipitated from the reaction solution by the addition of an aliphatic hydrocarbon solvent (e.g., hexane or heptane). The amount of the solvent to be used is 0.05 to 50 parts by weight, preferably 0.5 to 500 parts by weight per 1 part by weight of the solvent used for the preparation of the complex as above. After the addition of the solvent,
15 the resulting mixture may be cooled thereafter to a temperature of 0 to 30°C.

Said precipitated crystals of the chiral copper complex can be separated by filtration from the reaction mixture, and collected crystals may be further washed with said aliphatic or aromatic hydrocarbon solvent (e.g., an aromatic or aliphatic hydrocarbon solvent such as toluene, hexane,
20 heptane or the like), if necessary. The separated crystal can be subjected to high-performance liquid chromatography or elemental analysis to calculate the yield or purity thereof.

The chiral copper complex can form an adduct with a prochiral olefin of formula (5). Said adduct can be obtained by contacting the isolated chiral copper complex with the prochiral olefin of formula (5) or by using the
25 prochiral olefin as a solvent to produce the chiral copper complex. The adduct can be used as a catalyst in the cyclopropanation reaction.

The chiral copper complex having an optically active salicylideneaminoalcohol compound of formula (1) wherein R₁ and R₂ are the
30 same or different and independently represent an alkyl group, an aralkyl group, a phenyl group, a 2-methoxyphenyl group, a 2-tert-butoxy-5-tert-butylphenyl group or a 2-octyloxy-5-tert-butylphenyl group, and X₁ represents a nitro group and X₂ is a hydrogen atom, X₁ represents a chlorine atom and X₂ is a chlorine atom, or X₁ represents a
35 hydrogen atom and X₂ is a fluorine atom as shown by formula (1)' above can be obtained in a crystal form.

A description will be made to the process for producing optically active cyclopropanecarboxylic acid ester of formula (4) as defined above comprising reacting a prochiral olefin of formula (5) with a diazoacetic acid ester of formula (6) in the presence of a chiral copper complex thus obtained.

5 Examples of the alkoxy group represented by R_3 , R_4 , R_5 or R_6 of the prochiral olefin include an alkoxy group having 1 to 3 carbon atoms such as a methoxy, ethoxy, n-propoxy, i-propoxy, or the like.

 Examples of the alkyl group which may be substituted with a halogen atom or an alkoxy group include

10 a linear or branched alkyl group having 1 to 10 carbon atoms such as a methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, or n-decyl group,

 an alkyl group substituted with a halogen atom such as a fluorine atom, a chlorine atom, a bromine atom and the like a haloalkyl group such as
15 a chloromethyl, dichloromethyl, trichloromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trichloroethyl group or the like),

 an alkyl group substituted with an alkoxy group such as a methoxy, ethoxy, n-propoxy, i-propoxy group or the like.

 Examples of the alkylene group formed by R_3 and R_4 , or R_5 and R_6
20 include an alkylene group having 2 to 4 carbon atoms such as dimethylnene, trimethylene, or tetramethylene group.

 Examples of the alkenyl group which may be substituted with a halogen atom, an alkoxy group or an alkoxy carbonyl group, of which alkoxy may be substituted with a halogen atom or atoms, represented by R_3 , R_4 , R_5
25 include

 a linear or branched alkenyl groups having 1 to 10 carbon atoms such as an ethenyl, propenyl, 2-methylpropenyl, 1-butenyl, 2-butenyl, or hexenyl group,

 a haloalkenyl group, which is the above-described alkenyl group
30 substituted with the above-described halogen atom or atoms, such as a chloroethenyl group, a chloropropenyl group, 2,2-dichloroethenyl group, 2,2-difluoroethenyl group or the like,

 an alkoxy(C1-C3)carbonyl substituted alkenyl group such as
 2-methoxycarbonyl-2-methylethenyl group,
35 2-ethoxycarbonyl-2-methylethenyl group,
 2-(1,1,1,3,3,3-hexafluoroisopropoxycarbonyl)-1-methylethenyl group or the

like.

Specific examples of the prochiral olefin (5) include

- propene, 1-butene, isobutylene, 1-pentene, 1-hexene, 1-octene,
4-chloro-1-butene, 1-chloro-2-methylpropene, 2-pentene, 2-heptene,
5 2-methyl-2-butene, 2,5-dimethyl-2,4-hexadiene,
2-chloro-5-methyl-2,4-hexadiene, 2-fluoro-5-methyl-2,4-hexadiene,
1,1,1-trifluoro-5-methyl-2,4-hexadiene,
1,1,1-trifluoro-2-chloro-5-methyl-2,4-hexadiene,
2-methoxycarbonyl-5-methyl-2,4-hexadiene,
10 2-ethoxycarbonyl-5-methyl-2,4-hexadiene,
1,1-difluoro-4-methyl-1,3-pentadiene, 1,1-dichloro-4-methyl-1,3-pentadiene,
1,1-dichloro-2,4-dimethyl-1,3-pentadiene, 2-methyl-2,4-hexadiene,
2,3-dimethyl-2-pentene, 1,1,1-trichloro-4-methyl-3-pentene and the like.
1,1-dibromo-4-methyl-1,3-pentadiene,
15 1-chloro-1-fluoro-4-methyl-1,3-pentadiene,
1-chloro-1-(4-chlorophenyl)-4-methyl-1,3-pentadiene,
1-fluoro-1-bromo-4-methyl-1,3-pentadiene,
2-(1,1,1,3,3,3-hexafluoroisopropoxycarbonyl)-5-methyl-2,4-hexadiene,
1-methoxy-4-methyl-1,3-pentadiene,
20 1-ethoxy-4-methyl-1,3-pentadiene,
1-propoxy-4-methyl-1,3-pentadiene,
1-fluoro-1-methoxy-4-methyl-1,3-pentadiene,
1-fluoro-1-ethoxy-4-methyl-1,3-pentadiene,
1-fluoro-1-propoxy-4-methyl-1,3-pentadiene,
25 1,1,1-tribromo-4-methyl-3-pentene, 2-bromo-2,2-dimethyl-4-hexene,
2-chloro-2,5-dimethyl-4-hexene, 1-methoxy-2-methyl-1-propene,
1-ethoxy-2-methyl-1-propene, 1-propoxy-2-methyl-1-propene,
1-methoxy-8-methyl-2-butene, 1-ethoxy-3-methyl-2-butene,
1-propoxy-3-methyl-2-butene, 1,1-dimethoxy-3-methyl-2-butene,
30 1,1-diethoxy-3-methyl-2-butene, isopropylidenecyclopropane,
isopropylidenecyclobutane, 2-methyl-4-cyclopentylidene-2-buten,
isopropylidenecyclopentane and the like.

Preferably, R₃ and R₄, or R₅ and R₆ represent a methyl group. More preferred prochiral olefins are isobutylene and 2,5-dimethyl-2,4-hexadiene.

- 35 Examples of the alkyl group having 1 to 8 carbon atoms represented by R₇ in the diazoacetic acid ester of formula (6) include

a methyl group, an ethyl group, a n-propyl group, an isopropyl group, a n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a n-pentyl group, a n-hexyl group, a n-octyl group and the like.

5 Examples of the cycloalkyl group which may be optionally substituted with a lower alkyl group include a cyclohexyl group, a l-menthyl group, a d-menthyl group, an adamantyl group.

10 Examples of the phenyl group which may be optionally substituted with a lower alkyl group, a lower alkoxy group or a phenoxy group represented by R₇ in formula (6) include a phenyl group, 2-methylphenyl group, 3,5-dimethylphenyl group, 4-methyl-2,6-di-tert-butylphenyl group, 2-methoxyphenyl group, 3,5-dimethoxyphenyl group and the like.

15 Examples of the benzyl group which may be optionally substituted with a lower alkyl group, a lower alkoxy group, a phenoxy group or a halogen atom, represented by R₇ in formula (6) include a 3-phenoxybenzyl group, 2-methyl-3-phenylbenzyl group, 2,3,5,6-tetrafluorobenzyl group, 2,3,5,6-tetrafluoro-4-methylbenzyl group and the like.

20 Examples of the lower alkyl group which may be present on the cycloalkyl group or on the phenyl group include a (C1-C4)alkyl group such as a methyl group, an ethyl group, a n-propyl group, an i-propyl group, a n-butyl group, an i-butyl group, a sec-butyl group, and a t-butyl group.

Examples of the lower alkoxy group which may be present on the phenyl group include a (C1-C4)alkoxyl group such as a methoxy group, an ethoxy group, a n-propoxy group, an i-propoxy group, a n-butoxy group, an i-butoxy group, a sec-butoxy group, and a t-butoxy group.

25 Preferred are a (C1-C6)alkyl group,

a phenyl group which may be optionally substituted with a lower alkyl group, a lower alkoxy group or a phenoxy group such as a phenyl group, 2-methylphenyl group, 3,5-dimethylphenyl group, 4-methyl-2,6-di-tert-butylphenyl group, 2-methoxyphenyl group, 30 3,5-dimethoxyphenyl group,

a cyclohexyl group, a l-menthyl group, a d-menthyl group, and 2,3,5,6-tetrafluorobenzyl group and 3-phenoxybenzyl group.

Specific examples of the diazoacetic acid ester of formula (6) include ethyl diazoacetate, n-propyl diazoacetate, tert-butyl diazoacetate, phenyl 35 diazoacetate, l-menthyl diazoacetate, 4-methyl-2,6-di-tert-butylphenyl diazoacetate, 2,3,5,6-tetrafluorobenzyl diazoacetate, 3-phenoxybenzyl

diazoacetate and the like.

Said diazoacetic esters of formula (6) is commercially available or may be prepared by the known method such as a method of reacting corresponding amino acid esters with a diazotizing agent prepared from sodium nitrite and mineral acids may be used.

The reaction of prochiral olefin of formula (5) with diazoacetic esters of formula (6) in the presence of the prepared chiral copper complex catalyst is usually performed by adding the diazoacetic ester of formula (6) to a mixture of the copper complex catalyst and the prochiral olefin of (5) and optionally in a solvent. As described above, the present reaction may be performed in the presence of a reducing agent such as phenylhydrazine or the like.

An amount of prochiral olefin of formula (5) to be used is usually 1 or mole or more per mol of the diazoacetic acid ester of formula (6). The upper limit thereof is not particularly limited and, for example, a large excess amount may be used so as to serve as a reaction solvent.

An amount of the copper complex catalyst to be used is usually 0.001 to 5 mole %, preferably 0.03 to 1 mole % in terms of copper relative to the diazoacetic ester of formula (6).

Examples of the solvent to be used include a halogenated hydrocarbon such as 1,2-dichloroethane, chloroform, carbon tetrachloride or the like,

an aliphatic hydrocarbon such as hexane, heptane, cyclohexane and the like,

an aromatic hydrocarbon such as benzene, toluene, xylene and the like,

an ester such as methyl acetate, ethyl acetate and the like, and a mixture thereof. Alternatively, the prochiral olefin (5) may be used as a solvent.

An amount of a solvent to be used is usually 2 to 50 parts by weight, preferably 3 to 30 parts by weight per 1 part by weight of the diazoacetic ester (6).

A reaction temperature is usually -20 to 150°C , preferably -10 to 120°C . In addition, the reaction is usually performed under an inert gas atmosphere such as a nitrogen gas or the like and the inert gas may contain, for example, 5% or less oxygen gas. In addition, the reaction can be

conducted in the presence of excess amount of peroxide, for example, 20 moles or less, preferably 10 moles or less of peroxide per mol of the copper complex in the reaction without accompanying substantial adverse effect.

After completion of the reaction, the optically active
5 cyclopropanecarboxylic acid ester derivative of formula (4) can be separated by distillation or the like, which may be further subjected to ester hydrolysis or the like, or may be further purified, for example, by distillation, column chromatography or the like, if necessary.

Examples of the optically active cyclopropanecarboxylic acid esters of
10 formula (4) include optically active
methyl 2-methylcyclopropanecarboxylate,
ethyl 2-methylcyclopropanecarboxylate,
n-propyl 2-methylcyclopropanecarboxylate,
isopropyl 2-methylcyclopropanecarboxylate,
15 isobutyl 2-methylcyclopropanecarboxylate,
tert-butyl 2-methylcyclopropanecarboxylate,
cyclohexyl 2-methylcyclopropanecarboxylate,
menthyl 2-methylcyclopropanecarboxylate,
(4-methyl-2,6-di-tert-butylphenyl) 2-methylcyclopropanecarboxylate,
20 methyl 2,2-dimethylcyclopropanecarboxylate,
ethyl 2,2-dimethylcyclopropanecarboxylate,
n-propyl 2,2-dimethylcyclopropanecarboxylate,
isopropyl 2,2-dimethylcyclopropanecarboxylate,
isobutyl 2,2-dimethylcyclopropanecarboxylate,
25 tert-butyl 2,2-dimethylcyclopropanecarboxylate,
cyclohexyl 2,2-dimethylcyclopropanecarboxylate,
menthyl 2,2-dimethylcyclopropanecarboxylate,
(4-methyl-2,6-di-tert-butylphenyl) 2,2-dimethylcyclopropanecarboxylate,
methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,
30 ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,
n-propyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,
isopropyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,
isobutyl 2,2-dimethyl-3-(3-methyl-1-propenyl)cyclopropanecarboxylate,
tert-butyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,
35 cyclohexyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,
menthyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,

- (4-methyl-2,6-di-tert-butylphenyl) 2,2-dimethyl-3-(2-methyl-1-propenyl)-cyclopropanecarboxylate,
(3-phenoxybenzyl) 2,2-dimethyl-3-(2-methyl-1-propenyl)-cyclopropanecarboxylate,
5 (2,3,5,6-tetrafluoro-4-methoxymethylbenzyl) 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate,
methyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
ethyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
n-propyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
10 isopropyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
isobutyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
tert-butyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
cyclohexyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
menthyl 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
15 (4-methyl-2,6-di-tert-butylphenyl) 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)-cyclopropanecarboxylate,
(3-phenoxybenzyl) 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropane-carboxylate,
(2,3,5,6-tetrafluoro-4-methoxymethylbenzyl) 2,2-dimethyl-3-(2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
20 (2,2-dichloro-1-ethenyl)cyclopropanecarboxylate,
ethyl 2,2-dimethyl-3-(2,2-dibromo-1-ethenyl)cyclopropanecarboxylate,
methyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
ethyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
n-propyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
25 isopropyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
isobutyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
tert-butyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
cyclohexyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
menthyl 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
30 (4-methyl-2,6-di-tert-butylphenyl) 2,2-dimethyl-3-(2,2,2-trichloroethyl)-cyclopropanecarboxylate
(3-phenoxybenzyl) 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropane-carboxylate,
(2,3,5,6-tetrafluoro-4-methoxymethylbenzyl) 2,2-dimethyl-3-(2,2,2-trichloroethyl)cyclopropanecarboxylate,
35 (2,2,2-trichloroethyl)cyclopropanecarboxylate,
methyl 2,2-dimethyl-3-(2-carboethoxy-1-propenyl)cyclopropanecarboxylate,

- ethyl 2,2-dimethyl-3-(2-carbomethoxy-1-propenyl)cyclopropanecarboxylate,
n-propyl 2,2-dimethyl-3-(2-carboethoxy-1-propenyl)cyclopropanecarboxylate,
isopropyl 2,2-dimethyl-3-(2-carbomethoxy-1-propenyl)cyclopropane-
carboxylate,
5 isobutyl 2,2-dimethyl-3-(2-carboethoxy-1-propenyl)cyclopropanecarboxylate,
tert-butyl 2,2-dimethyl-3-(2-carbomethoxy-1-propenyl)cyclopropane-
carboxylate,
cyclohexyl 2,2-dimethyl-3-(2-carboethoxy-1-propenyl)cyclopropanecarboxylate,
menthyl 2,2-dimethyl-3-(2-carbomethoxy-1-propenyl)cyclopropane-
10 carboxylate,
4-methyl-2,6-di-tert-butylphenyl 2,2-dimethyl-3-(2-carboethoxy-
1-propenyl)cyclopropanecarboxylate
(3-phenoxybenzyl) 2,2-dimethyl-3-(2-carbomethoxy-1-propenyl)-
cyclopropanecarboxylate,
15 (2,3,5,6-tetrafluoro-4-methoxymethylbenzyl) 2,2-dimethyl-3-(2-carboethoxy-
1-propenyl)cyclopropane-carboxylate,
ethyl 2,2-dimethyl-3-(2-chloro-2-(4-chloropenyl)-1-ethenyl)cyclopropane-
carboxylate ,
ethyl 2,2-dimethyl-3-(2-chloro-2-trifluoromethyl-1-ethenyl)cyclopropane-
20 carboxylate and the like.

The optically active salicylideneaminoalcohol compound of formula (1)
contained in the residue after isolating the optically active
cyclopropanecarboxylic acid ester derivative of formula (4) can be recovered
by subjecting the residue to crystallization treatment, column
25 chromatography or the like.

Examples

The present invention will be illustrated by way of the following
Examples but are not to be construed to limit the present invention thereto.

30

Example 1

1.60 g of (R)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol, 0.98g of
2-hydroxy-5-nitrobenzaldehyde and 30 ml of toluene were mixed under a
nitrogen atmosphere, and the mixture was stirred at 80°C for 1 hour.
35 Thereafter, toluene was distilled off, followed by crystallization to obtain 1.35
g of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-(2-methoxyphenyl)-1-propanol.

Yield 74%

¹H-NMR (CDCl₃, TMS, ppm)

δ : 1.41 (d, 3H, J=6.6Hz), 3.62 (s, 6H), 5.26 (s, 1H), 5.66 (s, 1H),
5 6.65-7.27 (m, 8H), 7.52-7.67 (m, 2H), 8.03-8.10 (m, 3H)

Example 2

66.12 g of 2,5-dimethyl-2,4-hexadiene and 4 μg of phenylhydrazine were added to a 100 ml Schlenk tube purged with nitrogen, to which were
10 dropwise added a toluene solution of a copper complex prepared from 3.59mg of copper acetate monohydrate and 17.46 mg of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol, and a 9.9 g of a 2,5-dimethyl-2,4-hexadiene solution containing 2.28 g of ethyl diazoacetate over 2 hours at 80°C, respectively. Then, the mixture was stirred at the same temperature
15 for 30 minutes. Gas chromatography analysis of the resulting reaction mixture showed that the yield of ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was 97.4% and a trans/cis ratio was 59/41. It was shown by liquid chromatography analysis that optical purity of a (+)-trans-isomer was 56%ee and (+)-cis-isomer was 50%ee.

20 The reaction mixture was concentrated under reduced pressure to give a residue, which was analyzed by liquid chromatography. The recovery rate of the N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol remaining in the residue was 22% (relative to employed (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol).

25

Comparative Example 1

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was produced in 96.2% according to a similar manner as in Example 2 except that 14.60 mg of (R)-N-salicyliden-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was used instead of 17.46 mg of (R)-N-(5-nitrosalicylidene)-
30 2-amino-1,1-di(2-methoxyphenyl)-1-propanol. A trans/cis ratio was 60/40. The optical purity of a (+)-trans-isomer was 35%ee and the optical purity of a (+)-cis-isomer was 36%ee.

In addition, the recovery rate of the N-salicyliden-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was 5% (relative to the employed
35 N-salicyliden-2-amino-1,1-di(2-methoxyphenyl)-1-propanol).

Example 3

2.00 g of (R)-N-(3,5-dichlorosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was obtained by a similar manner as in
5 Example 1 except that 1.12 g of 2-hydroxy-3,5-dichlorobenzaldehyde was used instead of 0.98 g of 2-hydroxy-5-nitrobenzaldehyde. Yield 78%

¹H-NMR (CDCl₃, TMS, ppm)

δ : 1.35 (d, 3H, J=6.6Hz), 3.56 (d, 6H), 5.14 (q, 1H), 5.54 (s, 1H),
10 6.71-7.28 (m, 9H), 7.50-7.69 (m, 2H), 7.98 (s, 1H)

Example 4

2.0 g of (R)-2-amino-1,1-diphenyl-1-propanol, 1.47 g of 2-hydroxy-
5-nitrobenzaldehyde and 30 ml of methanol were mixed, and the mixture was
15 stirred at 80°C for 1 hour. Thereafter, the mixture was cooled to 20°C and the deposited crystals were collected by filtration. The collected crystals were washed with methanol and dried to give 3.22 g of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-diphenyl-1-propanol. Yield 97.2%

20 ¹H-NMR (CDCl₃, TMS, ppm)

δ : 1.29 (d, 3H, J=6.6Hz), 2.61 (s, 1H), 4.65 (q, 1H, J=6.6Hz), 6.82-6.89 (m, 1H),
7.20-7.54 (m, 10H), 8.12-8.15 (m, 2H), 8.26 (s, 1H)

Elemental analysis

25 Found : C 70.1%, H 5.4%, N 7.4%.
Calcd : C 70.2%, H 5.3%, N 7.5%.

Example 5

66.12 g of 2,5-dimethyl-2,4-hexadiene and 4 μg of phenylhydrazine
30 were added to a 100 ml Schlenk tube purged with a nitrogen gas having the oxygen concentration of 1%, and the mixture was stirred at 80°C for 30 minutes. Thereafter, a toluene solution of a copper complex catalyst prepared from 3.59 mg of copper acetate monohydrate and 8.73 mg of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol, and a
35 9.9 g of a 2,5-dimethyl-2,4-hexadiene solution containing 2.28 g of ethyl diazoacetate were added dropwise at the same temperature over 2 hours,

respectively. Thereafter, the mixture was further stirred and kept at the same temperature for 30 minutes, the resulting reaction mixture was analyzed by gas chromatography, which revealed that the yield of ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was 93.5% and a trans/cis ratio was 59/41. The optical purity of (+)-trans-isomer was 56%e.e. and 49%e.e. for a cis compound.

Comparative Example 2

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was produced according to a similar manner as that of Example 5 except that 7.83 mg of N-salicylidene-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was used instead of 8.73 mg of N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol. The yield was 89.3% and a trans/cis ratio was 60/40. The optical purity of (+)-trans-isomer was 34%ee and the optical purity of (+)-cis-isomer was 35%ee.

Example 6

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was produced according to a similar manner as that of Example 5 except that the reaction was performed in the presence of 0.066 g of cumene hydroperoxide which is an oxydizing agent and a 100 ml Schlenk tube purged with a nitrogen gas in Example 5. The yield was 95.8%, and trans/cis ratio was 58/42. The optical purity of (+)-trans-isomer was 61%ee and that of (+)-cis-isomer was 53%ee.

Comparative Example 3

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was produced according to a similar manner as that of Comparative Example 2 except that the reaction was performed in the presence of 0.066 g of cumene hydroperoxide which is an oxydizing agent and a 100 ml Schlenk tube purged with a nitrogen gas in Comparative Example 2. The yield was 94.2% and a trans/cis ratio was 60/40. The optical purity of a (+)-trans-isomer was 36%e.e. and that of (+)-cis-isomer was 36%e.e.

Example 7

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate

was produced according to a similar manner as that of Example 6 except that an amount of cumene hydroperoxide to be used was 0.26 g in Example 6. The yield was 81.1% and a trans/cis ratio was 59/41. The optical purity of (+)-trans-isomer was 46%e.e. and that of (+)-cis-isomer was 38%e.e.

5

Comparative Example 4

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was produced according to a similar manner as that of Comparative Example 3 except that an amount of cumene hydroperoxide to be used was 0.26g in Comparative Example 3. The yield was 8.5% and a trans/cis ratio was 59/41. The optical purity of (+)-trans-isomer was 21%e.e. and that of (+)-cis-isomer was 20%e.e.

10

Example 8

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was obtained in a similar manner as in Example 2 except that 5.22 mg of (R)-N-(3,5-dichlorosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was used instead of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol. The yield was 94.2% and a trans/cis ratio was 60/40. The optical purity of (+)-trans-isomer was 65%e.e. and that of (+)-cis-isomer was 60%e.e.

15

20

Example 9

66.12 g of 2,5-dimethyl-2,4-hexadiene and 4 µg of phenylhydrazine were added to a 100 ml Schlenk tube purged with nitrogen, a toluene solution of a copper complex catalyst prepared from 22.9 mg of a copper naphthenate/toluene solution containing 5 % by weight of copper metal copper and 7.53 mg of N-(5-nitrosalicylidene)-2-amino-1,1-diphenyl-1-propanol, and 9.9 g of a 2,5-dimethyl-2,4-hexadiene solution containing 2.28 g of ethyl diazoacetate were added dropwise at 80°C over 2 hours, respectively. Thereafter, the mixture was stirred and kept at the same temperature for 30 minutes, the resulting reaction mixture was analyzed by gas chromatography. The yield of ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was 96.3% and a trans/cis ratio was 61/39. The optical purity analyzed by liquid chromatography was 52%e.e. for (+)-trans-isomer and 44%e.e. for (+)-cis-isomer.

25

30

35

Example 10

3.8 g of (R)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol, 2.0 g of 2-hydroxy-3-fluorobenzaldehyde and 22 g of toluene were mixed and warmed to 80°C and stirred at the temperature for 1hr. Solvent was evaporated to give crystals, After 12 g of toluene were added thereto and warmed to dissolve the crystals at 80°C, 6 g of heptane were added thereto and cooled to a room temperature. Deposited crystals were collected by filtration, and washed with a mixed solvent of 3 g of toluene and 1.5 g of heptane, dried to give 4.37 g of (R)-N-(3-fluorosallylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol as yellowish powder. Yield: 80.7%. Mp. 116~118°C

¹H-NMR (CDCl₃, TMS, ppm)

δ : 1.34 (d, 3H), 3.56(d, 6H), 5.14(q, 1H), 5.49(s, 1H), 6.47-7.26(m, 9H), 7.56-7.68(m, 2H), 8.16(s, 1H).

Example 11

0.97 g of (R)-2-amino-1,1-di(2-n-butoxy-5-tert-butylphenyl)-1-propanol, and 0.33 g of 2-hydroxy-5-nitrobenzaldehyde were dissolved in a mixed solvent of 20 ml of ethanol and 20 ml of toluene, and refluxed for 1 hr, and the reaction mixture was evaporated to give 1.2 g of (R)-N-(5-nitrosallylidene)-2-amino-1,1-di(2-n-butoxy-5-tert-butylphenyl)-1-propanol as yellowish powder.

Yield: 94.9%. Mp. 67.9°C. [α]_D²⁵ -308° (c=0.1 % Chloroform).

¹H-NMR (CDCl₃, TMS, ppm)

δ : 0.87 - 0.93(m, 6H), 1.15(s, 9H), 1.29-1.31(m, 4H), 1.33(s, 9H), 1.36(s, 3H), 1.46 - 1.56(m, 4H), 3.72 - 3.85(m, 4H), 6.63-8.05(m, 9H), 8.57(s, 1H).

Elemental analysis

Found : C 71.5%, H 8.1%, N 4.6%.

Calcd : C 72.2%, H 8.2%, N 4.4%.

Example 12

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate was obtained in a similar manner as in Example 5 except that 9.2 mg of

(R)-N-(3,5-dichlorosalicyliden)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was used in place of 8.73 mg of (R)-N-(5-nitrosalicyliden)-2-amino-1,1-di(methoxyphenyl)-1-propanol.

Yield 94.4 %, Trans/cis ratio: 60/40, Optical purity of the trans-isomer: 58 %
5 e.e., Optical purity of cis-isomer: 53 % e.e.

Example 13

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropane-carboxylate was obtained in a similar manner as in Example 5 except that 8.19 mg of
10 (R)-N-(3-fluorosalicyliden)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was used in place of 8.73 mg of (R)-N-(5-nitrosalicyliden)-2-amino-1,1-di(methoxyphenyl)-1-propanol.

Yield 93.6 %, Trans/Cis ratio: 60/40, Optical purity of the trans-isomer: 46 %
e.e., Optical purity of Cis-isomer: 43 % e.e.

15

Example 14

Ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropane-carboxylate was obtained in a similar manner as in Example 7 except that 12.6 mg of
20 (R)-N-(5-nitrosalicyliden)-2-amino-1,1-di(5-tert-butyl-2-n-butoxyphenyl)-1-propanol was used in place of 8.73 mg of (R)-N-(5-nitrosalicyliden)-2-amino-1,1-di(methoxyphenyl)-1-propanol.

Yield 91.3 %, Trans/cis ratio: 56/44, Optical purity of the trans-isomer: 51 %
e.e. Optical purity of cis-isomer: 35 % e.e.

25

Example 15

0.52 g of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(5-tert-butyl-2-n-butoxyphenyl)-1-propanol prepared in Example 11, 0.164 g of copper acetate monohydrate and 40 ml of toluene were charged in a 100 ml glass flask, and the resulting mixture was reacted at 80°C for 1 hr. After cooled to a room
30 temperature, the reaction solution was transferred to a separatory funnel and thoroughly mixed with 30 ml of an aqueous sodium hydroxide solution, then settled and separated. Separated layer was washed twice with deionized water and separated organic phase was evaporated to give 0.53 g of copper complex as green powder.

35 Decomposition Temp : 110 - 130°C $[\alpha]_{546} +314^{\circ}$ (c=0.1% chloroform)
Elemental Analysis

Found: C 66.9%, H 7.4 %, N 3.8 %, Cu 7.66 %.

Example 16

6.9 mg of the copper complex prepared in Example 15, 10 ml of ethyl acetate and 1.1 mg of phenylhydrazine were charged in a 50 ml Schlenk glass tube reactor purged with nitrogen gas equipped with a magnetic stirrer to give a catalyst solution. 5 ml of the catalyst solution and 3.5 g of isobutylene were charged in a 100 ml stainless steel autoclave purged with nitrogen gas, and to the resulting solution was added 10 ml toluene solution containing ethyl diazoacetate, of which content is 20 mmol, at 40°C for 2 hrs. After stirring at the same temperature for 30 min, the reaction solution was cooled. Content of ethyl 2,2-dimethyl-3-(2-methyl-1-propenyl)-cyclopropanecarboxylate was analyzed by gas chromatography analysis. Yield: 91%.

After hydrolysis of the obtained ester, the acid was derivatized to its 1-menthyl ester and the optical purity was measured to show 87 % e.e.

Example 17

9.8 g of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol, 4.0 g of copper acetate monohydrate, and 550 g of toluene were charged in a 100 ml glass flask and reacted at 80°C for 1 hr. After being cooled to a room temperature, 8.7 g of 28 (wt)% sodium methylate were added thereto and stirred for 10 min. Then the reaction mixture was mixed with 200 g of water and separated by a separatory funnel. Separated oil layer was evaporated to give 11.2 g of copper complex as green powder.

Decomposition Temp : 158~165°C. $[\alpha]_{546} +927^{\circ}$ (c=0.1%, Chloroform)

Elemental Analysis

Found : C 60.1%, H 4.9%, N 5.2%, Cu 10.4%

Example 18

0.5 g of the copper complex obtained in Example 17, 100 g of 2,5-dimethyl-2,4-hexadiene were charged in a 100 ml glass flask, and reacted at 80°C for 1 hr under stirring. A green solid was precipitated on cooling the reaction mixture. Precipitated crystals were collected by filtration and dried to give 0.4 g of an adduct as green powder.

Elemental Analysis

Found : C 61.1 %, H 5.3 %, N 5.0%, Cu 11.7%.

Calcd: C 60.8 %, H 5.3 %, N 5.1 %, Cu 11.5 %.

- 5 (Calcd as an adduct of two molecules of a copper complex and one molecule of 2,5-dimethyl-2,4-hexadiene)

Example 19

- 5.53mg of the adduct obtained in Example 18, 66.1 g of
10 2,5-dimethyl-2,4-hexadiene, and 4 μ g of phenylhydrazine were charged in a Schlenk tube purged with nitrogen gas, to the resulting mixture was dropwise added a 9.9 g of a solution of 2,5-dimethyl-2,4-hexadiene solution containing 2.3 g of ethyl diazoacetate over 2 hrs and kept at the same temperature for 30 min under stirring. The resulting reaction mixture was
15 analyzed by gas-chromatography analysis. 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropane-carboxylate was obtained in 97.2 % yield. Trans/cis ratio was 58/42. Optical purities of the trans-isomer and cis-isomer were 62.5% e.e. and 53.2 % e.e.

20 Example 20

- 19.6 g (44.9 mmol) of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol, 8.96 g (44.9 mmol) of cupric acetate, and 160 g of toluene were mixed in a flask and reacted at 80°C for 1 hr under stirring. The reaction solution was cooled to 10°C and blue-green chrystalls
25 were deposited. Deposited crystals were collected by filtration, washed with 50 g of cold toluene, and dried at room temperature to give 19.1 g of a copper complex of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol in a yield of 82.0%.

30 Example 21

- 19.6 g (44.9 mmol) of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol, 8.96 g (44.9 mmol) of cupric acetate, and 160 g of toluene were mixed in a flask and reacted at 80°C for 1 hr under stirring. 100 g of n-heptane were added to the reaction solution to precipitate
35 blue-green chrystalls. The precipitated reaction mixture was cooled to 10°C and filtered to collect the crystals. Collected crystals were washed with 100 g

of n-heptane, and dried at room temperature to give 22.1 g of a copper complex of (R)-N-(5-nitro-salicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol in a yield of 99.0%.

5 Example 22

0.46 of a copper complex of (R)-N-(3,5-dichlorosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was obtained in a yield of 97.9% according to a similar manner as in Example 21 except that 0.415 g (0.901 mmol) of (R)-N-(3,5-dichlorosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was used in place of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol and 0.180 g (0.901 mmol) of cupric acetate, 10 g of toluene, 10 g of n-heptane for precipitation of the crystals, 10 g of n-heptane for washing the crystals were used.

15 Example 23

0.434 of a copper complex of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-diphenyl-1-propanol was obtained in a yield of 99.1 % according to a similar manner as in Example 21 except that 0.376 g (1.00 mmol) of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-diphenyl-1-propanol was used in place of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol and 0.20g (1.0 mmol) of cupric acetate, 10 g of toluene, 10 g of n-heptane as a poor solubility solvent, 10 g of n-heptane for washing the crystals were used.

 Example 24

25 33.06 g (300mmol) of 2,5-dimethyl-2,4-hexadiene, 4.97 mg (0.01 mmol) of a copper complex of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol were charged in a 100 ml Schlenk tube purged with nitrogen gas and 4 μ g of phenylhydrazine was added thereto. 1.14 g (10 mmol) of ethyl diazoacetate was added the resulting solution over 2
30 hrs and stirred at the same temperature for 30 min. The gas chromatography analysis of the reaction solution showed that the yield of chrysanthemate was 97.6 %, trans/cis ratio was 58/42. High-performance liquid chromatography analysis showed that optical purity of the trans-isomer was 63 % e.e and cis-isomer was 57 % e.e.

35

 Example 25

Chrysanthemate was produced in a similar manner as in Example 24 in a yield of 97.6 %, where trans-cis ratio was 60/40, trans-isomer was 61 % e.e and cis-isomer was 56 % e.e. except that 5.22 mg (0.01 mmol) of a copper complex of (R)-N-(3,5-dichlorosalicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol was used in place of 4.97mg of a copper complex of (R)-N-(5-nitro-salicylidene)-2-amino-1,1-di(2-methoxyphenyl)-1-propanol.

Comparative Example 5

1.0 g (2.56 mmol) of (R)-N-(5-nitrosalicylidene)-2-amino-1,1-diphenyl-1-propanol and 0.511 g (2.56 mmol) of cupric acetate were mixed in 5 g of toluene and reacted at 80°C for 1 hr under stirring. Then 50 g of n-heptane was added thereto and cooled to 10°C, which produced no precipitated product and remain as a clear solution.

(R)-N-(5-nitrosalicylidene)-2-amino-1,1-diphenyl-1-propanol